

GIARDIA IN DOGS

What are Giardia?

Giardia are sometimes confused with worms because they invade the gastrointestinal tract and can cause diarrhea. They are not worms; instead they are one-celled parasites classified as protozoa.

Most dogs that are infected with *Giardia* do not have diarrhea or any other signs of illness. When the eggs (cysts) are found in the stool of a dog without diarrhea, they are generally considered a transient, insignificant finding. However, in puppies and debilitated adult dogs, they may cause severe watery diarrhea that may be fatal.

How did my dog get Giardia?

A dog becomes infected with *Giardia* when it swallows the cyst stage of the parasite. Once inside the dog's intestine, the cyst goes through several stages of maturation. Eventually, the dog is able to pass the infective cysts in the stool. These cysts lie in the environment and can infect other dogs. They may also be transmitted through drinking *infected* water.

How is giardiasis diagnosed?

Giardiasis is diagnosed by performing a microscopic examination of a stool sample. The cysts are quite small and usually require a special flotation medium for detection, so they are not normally found on routine fecal examinations. Occasionally, the parasites may be seen on a direct smear of the feces. A blood test is also available for detection of antigens (cell proteins) of *Giardia* in the blood. This test is probably more accurate than the stool exam, but it requires several days to get a result from the laboratory performing the test.

How is giardiasis treated?

The typical drugs used to kill *Giardia* are Metronidazole and Fenbendazole, two antibiotic-type drugs. They are usually given for 5 to 7 days, then repeated a few weeks later. Other drugs are also used if diarrhea and dehydration occur. Because cysts can stick to the fur of the infected patient and be a source for re-infection, the positive animal should receive a bath at least twice in the course of treatment.

Can humans become infected with Giardia?

Giardia can also cause diarrhea in humans. Therefore, environmental disinfection is important. The use of chlorine bleach, one cup per gallon (250 ml per 4 liters) of water is effective if the surfaces and premises can be safely treated with it.